

# Proceedings of the Iowa Academy of Science

---

Volume 35 | Annual Issue

Article 49

---

1928

## X-Ray Diffraction in Liquids: Isomers of Normal Heptane

G. W. Stewart

Copyright © Copyright 1928 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Stewart, G. W. (1928) "X-Ray Diffraction in Liquids: Isomers of Normal Heptane," *Proceedings of the Iowa Academy of Science*, 35(1), 250-250.

Available at: <https://scholarworks.uni.edu/pias/vol35/iss1/49>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## X-RAY DIFFRACTION IN LIQUIDS: ISOMERS OF NORMAL HEPTANE

G. W. STEWART

The following isomers of normal heptane were kindly supplied by the Ethyl Gasoline Corporation:

3-3, 2-2, 2-3 and 2-4 dimethyl pentane, 3 ethyl pentane, 3 methyl hexane, 2 methyl hexane, 2-2-3 trimethyl butane.

The conclusions obtained from x-ray diffraction measurements are as follows:

1. Dimethyl pentane, whether 2-3, 2-4, 2-2 or 3-3, has a greater diameter than normal heptane by  $1.04\text{\AA}$  or about twice the increase caused by two such branches in tertiary butyl alcohol, and roughly twice the increase caused by a branch  $\text{CH}_3$  or of OH in the case of the alcohols.

2. The increase in diameter with the two  $\text{CH}_3$  groups attached either to adjacent or nonadjacent C atoms seems to alter the diameter by the same amount,  $1.04\text{\AA}$ . This is shown by the four heptanes just enumerated.

3. The attachment of two  $\text{CH}_3$  groups to the same atom may or may not increase the diameter by the same amount,  $1.04\text{\AA}$ , this depending upon whether a third  $\text{CH}_3$  branch is attached to an adjacent atom (2, 2, 3 trimethyl butane or 2, 2, 4 trimethyl pentane).

4. The union of three like branches, as is 3 ethyl pentane increased the diameter by an amount indicating the probable breaking up of the "straight" chain into a symmetrical form with equal angles between each pair.

5. The area of cross-section of the molecule when the structure changes to an isomer decreases much more rapidly than computations indicate that it should. This suggests a change in the structure of the chain.

6. The paraffin molecule is a chain that is similar to but not like the alcohol chain, in that the attachment of side branches does not indicate equal alterations in diameter.

7. The structure of 3 methyl hexane is better termed 2 ethyl pentane.

8. The attachment of a  $\text{CH}_3$  group on the next to the end C atom evidently alters the end of the chain, causing a symmetrical arrangement of the two  $\text{CH}_3$  groups and increasing the diameter of the molecule by a relatively small amount.